



MOOIVET

Animal Health Products

— Est. 1985 —

AGITA™
WG 10

Dosage form: Paint-on.

Presentation: 50 g tub and 5 kg.

Composition: Thiamethoxam (neonicotinoid) 100 g/kg, (Z) 9-Tricosene 0,50 g/kg.

Storage instructions: Store below 30 °C.

Indication:

Adult fly control. A water-dispersable paint-on residual action stomach and contact insecticide with an attractant for the control of flies and litter beetles (*Alphitobius diaperinus*) in poultry houses.

Warnings:

- Store in original container and do not keep or apply where food or feed may become contaminated.
- Poisonous by contact and when ingested.
- Toxic to bees and aquatic organisms.
- Keep out of reach of children, uninformed persons and animals.
- Do not store the made-up liquid because it will lose its efficacy.
- Keep container tightly closed between use to maintain strength of attractant.

Precautions:

- Wear rubber gloves when applying. In case of accidental contact, remove contaminated clothing and wash skin thoroughly with soap and water.
- Do not eat, drink or smoke during application.
- Destroy empty container by perforation and flattening.
- Never re-use empty container for any other purpose.
- Prevent contamination of food, feedstuffs, eating utensils and drinking water. Clean application equipment after use and do not contaminate crops, grazing, rivers, dams and areas not under treatment with wash water.

Directions for use: Use only as directed.

AGITA™ WG 10 Fly Bait is suitable for use in stables, poultry houses, pigsties, dairies

and other farm buildings as well as industrial buildings where flies are a nuisance and a health hazard. **AGITA™ WG 10** Fly Bait will also control litter beetles (*Alphitobius diaperinus*) infestations in broiler houses.

As a paint-on for adult fly control:

Thoroughly mix 50 g **AGITA™ WG 10** Fly Bait with 40 ml lukewarm water until a brushable liquid is obtained. Sufficient for a building with a ground area of 20 m² or corresponding wall or ceiling surface of 40-60 m². Stir regularly. Surfaces to be treated should be as dust-free as possible. Apply in strips of about 30 cm² to the walls and ceiling where flies congregate, e.g. sunny areas, window frames, supporting posts, etc. Cover about 2,5 % of the total wall and ceiling surface areas. Should it be unacceptable to treat surfaces in this way, strips of white carton or plastic may be painted and suspended from the roof. Stir regularly. Repeat treatment as necessary.

As a band spray or feed bait for litter beetle control in poultry houses:

(Claim printed on 5 kg packs only.)

Band spray: The following application method for **AGITA™ WG 10** Fly Bait is recommended:

1. The application rate is 400 mg of active ingredient (4 g of **AGITA™ WG 10** Fly Bait) per m² of surface area.
2. The product must be mixed into sufficient water to allow a thorough wetting of a one metre-wide band on all sides of the house (+/-1 litre solution per 15 m²).
3. The recommended position of the band spray is at the junction of the wall and ceiling, with one-third of the band on the wall and two-thirds on the ceiling. Furthermore, the area around any cracks or openings in the ceiling should also be sprayed. The rationale



for this programme is the fact that the beetles hide in the ceiling between cycles and then crawl out just before the new chicks arrive. At this point, therefore, they will come into contact with the spray band.

4. The spray should be applied after cleaning and the placement of new litter, just before the house is heated up for the arrival of the new batch of chicks.

Feed bait: The following method of application of **AGITA™ WG 10** Fly Bait is recommended:

1. Mix 25 g of **AGITA™ WG 10** Fly Bait with 1 kg of feed. Place 30 g of this mixture under each feeder pan. On an average house of 500 pans and placing 30 g of mixture under each pan a house would require 375 g of **AGITA™** in 15 kg of feed. If chick paper is used, put the bait down before the chick paper is placed under the feeders.
2. If required, a second application can be done midway through the cycle.
3. The safety profile of the product is such that any small amounts of **AGITA™ WG 10** Fly Bait that may be consumed by the chickens will not pose any health threat to the chickens nor the consumer.
4. The rationale for this method of treatment is the fact that the beetles and larvae will congregate around the feeder pans which provide a source of food for them.

Agri Farmacy SA (Pty) Ltd. | Reg.: 2020/095840/07

96 Dr. James Moroka Avenue, Potchefstroom, 2531 | P.O. Box 1768, Potchefstroom, 2520

E-mail: info@agrifarmacysa.co.za | Tel.: +27 18 297 8155

www.agrifarmacysa.co.za

Product Reg. No: L6670 (Act 36/1947)

Distributed by: MOOIVET VETERINARY PRODUCTS CC (Co. Reg. No.: 1985/010721/23)

Registration holder: ELANCO ANIMAL HEALTH, N1 Business Park, Old Pretoria Road, Kosmosdal Ext 7, 0157





MOOIVET

Animal Health Products

— Est. 1985 —



AGITA[™]
WG 10

Agri Pharmacy SA (Pty) Ltd. | Reg.: 2020/095840/07
96 Dr. James Moroka Avenue, Potchefstroom, 2531 | P.O. Box 1768, Potchefstroom, 2520
E-mail: info@agrifarmacysa.co.za | Tel.: +27 18 297 8155

www.agrifarmacysa.co.za

Product Reg. No: G0641 (Act 36/1947)
Distributed by: MOOIVET VETERINARY PRODUCTS CC (Co. Reg. No.: 1985/010721/23)
Registration holder: ELANCO ANIMAL HEALTH, N1 Business Park, Old Pretoria Road, Kosmosdal Ext 7, 0157

