



MOOIVET

Animal Health Products

— Est. 1985 —

ECTOMIN™ 100 EC

Dosage form: Dip and spray.

Presentation: 1 L, 5 L, 20 L.

Composition: Cypermethrin (high-cis) (pyrethroid) 100 g/L.

Indication: Store in tightly closed original container in a cool, dry place away from direct sunlight.

Warnings:

- No withdrawal period for milk, meat or eggs.
- Handle with care.
- Poisonous by contact, swallowing or inhalation.
- Store concentrate in original container away from food and feed.
- Keep out of reach of uninformed persons, children and pets.
- Toxic to fish, bees and reptiles.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions:

- Do not mix **Ectomin™** with any other dip.
- Avoid contact with concentrate. In case of accidental eye contact, flush eyes with plenty of running water for at least 15 minutes. In case of accidental skin contact, remove contaminated clothing and wash skin thoroughly with soap and water.
- Avoid prolonged contact with dipwash or inhalation of spray mist.
- Do not eat, drink or smoke while mixing or during use of dip.
- Prevent contamination of food, food utensils, grazing, drinking water, dams, rivers, etc.

Directions for use: Directions for use: Use

only as directed. Shake well before use.

Premix by adding the measured amount of concentrate slowly to double the volume of water whilst stirring. After thoroughly stirring this mixture, add it to the bulk of the water in the diptank or spray race sump. Make sure that in the case of plunge tanks, the larger volume is poured in at the deep end. Mix contents of the tank well before commencing dipping or spraying.

Recommendations:

Spray races: Mix only enough spray-wash for one treatment and do not allow wash to stand over to the next spraying.

Boosting: Boosting of the spray-wash is accomplished by adding 200 ml **Ectomin™** after every 100 head of cattle.

Hand spraying: Thoroughly wet the animal to the skin, especially under the tail, the ears and between the legs where ticks cluster.

Dipping intervals:

Ticks, flies and keds: Dip or spray weekly.

Lice: Dip or spray when necessary.

In areas with a high *Amblyomma* tick challenge, it is recommended that

Ektoban™ (Reg. No. G0598 Act 36/1947) is used.

Dilution rates:

	FRESHFIL		REPLENISH	
	ECTOMIN™	WATER	ECTOMIN™	WATER
Plunge-drips	1 L	1000 L	1 L	1000L
Spray race	1 L	1000 L	1 L	1000L
Hand-spraying: 10 ml Ectomin™ per 10 L water				

For the control of lice on sheep and goats, use 2 ml Ectomin™ / 10 L water applied once only. For the control of poultry mites (*Dermanyssus gallinae* and *Ornithonyssus sylviarum*), dilute 60 ml **Ectomin™** in 60 L of water, sufficient to treat 1000 birds.



Agri Farmacy SA (Pty) Ltd. | Reg.: 2020/095840/07
96 Dr. James Moroka Avenue, Potchefstroom, 2531 | P.O. Box 1768, Potchefstroom, 2520
E-mail: info@agrifarmacysa.co.za | Tel.: +27 18 297 8155

www.agrifarmacysa.co.za

Product Reg. No: G3314 (Act 36/1947)
Distributed by: MOOIVET VETERINARY PRODUCTS CC (Co. Reg. No.: 1985/010721/23)
Registration holder: ELANCO ANIMAL HEALTH, N1 Business Park, Old Pretoria Road, Kosmosdal Ext 7, 0157

