



MOOIVET⁺

Animal Health Products

— Est. 1985 —



Dosage form: Water-soluble granules.

Presentation: 4 kg, 20 kg.

Composition: Cyromazine (triazine) 20 g/kg.

Storage instructions: Store below 30 °C in original container and in a cool place. Protect from direct sunlight and humidity.

Indication: Water-soluble, granular larvicide that prevents fly and mosquito larval development in breeding sites.

Warnings:

- Take care not to overdose when applying to water.
- Harmful if ingested.
- Harmful to aquatic life with long lasting effects.
- Avoid release into the environment.
- Store in original container in a cool place away from food and feedstuffs.
- Keep out of the reach of uninformed persons, children and animals.
- Do not store made up product overnight as it will lose its efficacy.
- Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the remedy concerned as well as the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the

label instructions or to the occur

Precautions:

- Wear overalls and rubber gloves when applying.
- In case of accidental contact, remove contaminated clothing and wash skin thoroughly with soap and water.
- If medical advice is needed, have the product container or label at hand.
- Do not eat, drink or smoke whilst mixing or applying Neporex™ before having washed with soap and water.
- Destroy empty container by perforation and flattening and never re-use for any other purpose.
- Prevent contamination of food, feedstuffs and drinking water.
- Do not use in slurry intended for biogas production.
- Dispose of contents/container to an approved incineration plant.

Directions for use: Use only as directed.

Compatibility: Do not mix with any other insecticide.

Neporex® prevents the development of fly and mosquito larvae in breeding sites. It has no effect on adult insects and the reduction in adult numbers is therefore gradual. **Neporex™** must be applied as early in the breeding season as possible to avoid the build-up of large numbers of adult insects.

- **Flies:** **Neporex™** is applied by spreading, sprinkling, or spraying onto the fly breeding sites. Treat manure, refuse and compost heaps, dung pits, slurry and dung channels or soiled litter. Apply treatment at 14-day intervals until control is achieved. Where adult flies from outside migrate into animal units, adult fly control should be practised.
- **Poultry units with dung pits:** Treat the whole dung area evenly. First

treatment: Ideally when the dung is less than 10 cm deep.

Second treatment: Fortnight later. A third treatment may be necessary if the first treatment was made on deep dung.

- **Pig units with slatted floors:** Fully slatted floors: Treat the whole floor area. For all-in-all-out systems, treat empty unit between batches. Other systems: treat when necessary.
- Partly slatted floors: Treat whole of slatted area and other areas where dung and waste accumulate.
- Manure and refuse heaps: Treat regularly as new material is added.
- Calf-fattening, dairies, feedlots and horse stables: The fly breeding sites are to be found along the walls and around and under installations, where the bedding and manure are not compacted by the animals. Treat approximately 50 cm wide strips evenly in these sites. Neporex™ needed for 10 m² of surface area:
 - Wet surfaces: 250 g dry granules scattered.
 - Dry surfaces: Watering can – 250 g dissolved into 10 L water; sprayer – 250 g dissolved into 1 to 4 L of water.
 - Mosquitoes: Treat all breeding sites, including ponds, stagnant pools, swamps, marshes, puddles and other standing water. Sprinkle Neporex™ granules evenly over the water surface at a rate of 50 g/1000 L water. Repeat treatment every 8 weeks or at the first sign of breeding activity.

1000 L water is equivalent to:

WATER DEPTH (CM)	10	20	30	40	50	75	100
WATER SURFACE (M ²)	10	5.0	3.3	2.5	2.0	1.3	1.0

Agri Farmacy SA (Pty) Ltd. | Reg.: 2020/095840/07
 96 Dr. James Moroka Avenue, Potchefstroom, 2531 | P.O. Box 1768, Potchefstroom, 2520
 E-mail: info@agrifarmacysa.co.za | Tel.: +27 18 297 8155

www.agrifarmacysa.co.za

Product Reg. No: L3055 (Act 36/1947)
 Distributed by: MOOIVET VETERINARY PRODUCTS CC (Co. Reg. No.: 1985/010721/23)
 Registration holder: ELANCO ANIMAL HEALTH, N1 Business Park, Old Pretoria Road, Kosmosdal Ext 7, 0157

